Non-Invasive Prenatal Testing – NIPT

NIPT screens a maternal blood sample for chromosomal aneuploidy in fetal DNA using the following methodology:

- (1) Extraction of fetal cell-free DNA from the maternal blood sample
- (2) High throughput sequencing of the extracted fetal cell-free DNA
- (3) Calculation of molecular mass of fetal DNA in all chromosomes

Based on the scope, NIPT can screen the following conditions:

- (a) Whole Genome 23 pairs of human chromosomes
- (b) Common Chromosomal abnormality:

Trisomy 13 (Patau's Syndrome)

Trisomy 18 (Edwards' Syndrome)

Trisomy 21 (Down's Syndrome)

NIPT is capable of genome-wide aneuploidy detection of the whole fetal genome (23 pairs of chromosomes). Test results with the interpretation of risk for Trisomy 13 Trisomy 18, Trisomy 21 and sex chromosome aneuplodies will be provided. This test confers an accuracy of up to **99%** on the detection of fetal chromosomal aneuploidy.

Pregnancy Type	Blood ID	Patient ID
Singleton	NA	NA

Test Results Summary

Autosomal Aneuploidies		Risk	Test Results	Aneuploidy Risk	
Chromosome 13			Low risk group	< 1/10000	
Chromosome 18			Low risk group	< 1/10000	
Chromosome 21			Low risk group	< 1/10000	
Other Chromosomes			Low risk group	-	

Dr. Parag Tamhankar DM (Medical Genetics), MD, DNB (Pediatrics) Senior Genetic Consultant Dr.Nirmal A. Vaniawala MD (Path. & Bact.)

Mariawala

Test Results for Sex Chromosome Aneuploidies

Sex Chromosome Aneuploidies	Risk	Test Results	Aneuploidy Risk
хо		Low risk group	< 1/10000
XXY		Low risk group	< 1/10000
XXX		Low risk group	< 1/10000
XYY		Low risk group	< 1/10000

※ Risk description:

■ Low risk group;

■ Borderline group;

● High risk group

Test Results for Other Chromosomal Aneuploidies

Other Chrom	osomal Aneuploidies	Risk	Test Results
Chromosome 1			Low risk group
Chromosome 2			Low risk group
Chromosome 3			Low risk group
Chromosome 4			Low risk group
Chromosome 5			Low risk group
Chromosome 6			Low risk group
Chromosome 7			Low risk group
Chromosome 8			Low risk group
Chromosome 9			Low risk group
Chromosome 10			Low risk group
Chromosome 11			Low risk group

Dr. Parag Tamhankar DM (Medical Genetics), MD, DNB (Pediatrics) Senior Genetic Consultant Dr.Nirmal A. Vaniawala MD (Path. & Bact.)

Chromosome 12		Low risk group		
Chromosome 14		Low risk group		
Chromosome 15		Low risk group		
Chromosome 16		Low risk group		
Chromosome 17		Low risk group		
Chromosome 19		Low risk group		
Chromosome 20		Low risk group		
Chromosome 22		Low risk group		
※ Risk description: Low risk group; Borderline group, High risk group				

Sample information


Fetal DNA fraction


10.09%


Note: In rare cases when fetal DNA fraction level is low, new blood sample will be requested for retesting.


Dr. Parag Tamhankar DM (Medical Genetics), MD, DNB (Pediatrics) Senior Genetic Consultant Dr.Nirmal A. Vaniawala MD (Path. & Bact.)

Test Results










• These images are only for representation puposes.

Dr. Parag Tamhankar DM (Medical Genetics), MD, DNB (Pediatrics) Senior Genetic Consultant

Maniewala

Dr.Nirmal A. Vaniawala MD (Path. & Bact.)

Prenatal Examination



About the Test

YES-IN-GENE specializes in research development and high-quality professional services in clinical genetic testing. NIPT analyzes circulating fetal cell-free DNA extracted from a maternal blood sample, and is offered to pregnant women with a pre-test risk of aneuploidy in chromosomes such as 13, 18, 21, X or Y. The chance that a fetus is affected with chromosomal aneuploidy can be estimated using bioinformatics analyses, by which the accuracy rate and sensitivity are over 99%. The accuracy and quality of the test may be affected by low fetal fraction, high data noise due to improper blood sample collection, handling, storage, or transportation.

Limitations of the Test

Non-invasive prenatal testing should only be considered a screening test. The screening test of fetal cell-free DNA cannot compare with the prenatal diagnosis with Amniocentesis or Chorionic Villus Sampling (CVS). Pregnant women with a positive NIPT screening result should be given an invasive prenatal diagnosis and referred further for genetic counselling to confirm conditions. On the other hand, a negative test result does not ensure an unaffected pregnancy. Even though NIPT provides reliable results, it does not apply to all cases of chromosomal abnormalities, for example, cases due to placental, maternal, or fetal mosaicism, or other causes (e.g. micro-deletions, chromosome re-arrangements, translocations, inversions, unbalanced translocations, uniparental disomy, etc.). NIPT is also not applicable for cases with a diagnosed multiple gestation, or with gestational age that is less than 10 weeks. In rare cases when a borderline screening result is reported, retesting is required to confirm conditions.

Dr. Parag Tamhankar DM (Medical Genetics), MD, DNB (Pediatrics) Senior Genetic Consultant Dr.Nirmal A. Vaniawala MD (Path. & Bact.)

Test Method

NIPT applies a non-invasive and low-risk procedure to collect fetal DNA samples. Circulating fetal cell-free DNA is purified from the plasma component of 10mL anti-coagulated maternal whole blood. It is then converted into a genomic DNA library for Next Generation Sequencing to determine Trisomy 21, 18 and 13 and other chromosomal abnormalities.

References

- 1. Obstet Gynecol 2012; 119:890-901.
- 2. BMJ 2011; 342:c7401.
- 3. Prenat Diagn 2012; 32:c7401.
- 4. ACOG/SMFM Joint Committee Opinion No. 545, Dec 2012.

Note:	The sex	of fetus	s is not	revealed	due	to PND	T Act.
		E	End Of	Report			

Dr. Parag Tamhankar DM (Medical Genetics), MD, DNB (Pediatrics) Senior Genetic Consultant Dr.Nirmal A. Vaniawala MD (Path. & Bact.)